

Effect of streptozotocin-induced diabetes on kidney Na^+/K^+ -ATPase¹

Ágota VÉR, Ildikó Szántó, P. Csermely, K. Kalff, Edit Végh,
T. Bányász*, Z. Marcsek**, T. Kovács*, J. Somogyi

Institute of Biochemistry I, Semmelweis University of Medicine, Budapest

* Department of Physiology, University Medical School of Debrecen, and

** Joint Research Organization of Hungarian Academy of Sciences, Budapest, Hungary

Received April 10, 1995

Accepted May 30, 1995

The maximal capacity of low affinity ouabain binding sites in kidney medulla was found to be increased by $20 \pm 3.8\%$ after 2 weeks, and by $35 \pm 4.5\%$ in 4 weeks diabetes. However, in kidney cortex no similar changes could be detected. Western blot analysis of Na^+/K^+ -ATPase subunits in kidney medulla indicated a significant enhancement of both the α_1 and β_1 subunit in two and four weeks diabetic rats (α_1 : 35 ± 3.1 , $51 \pm 5.8\%$ and β_1 : 31.3 ± 5.2 and $43.2 \pm 6.8\%$, respectively). However, kidney cortex showed no significant change in any condition tested. In diabetes we could detect a significant change only in the medulla in case of the b subunit mRNA transcript, which showed 1.69 ± 0.59 and 2.89 ± 0.81 times increased in two and four weeks diabetic state, respectively. There was no change in the α_1 subunit mRNA abundance. Insulin treatment of diabetic animals did not result in a complete reversal of diabetes-induced changes in ouabain binding capacity or in the amount of Na^+/K^+ -ATPase α_1 and β_1 subunit protein and mRNA levels. Our data indicate a good correlation between changes in low affinity ouabain binding capacity and the level of α_1 isoform in diabetic rats, and suggest an important role of the b subunit in the regulation of enzyme quantity.

Keywords: streptozotocin-induced diabetes, Na^+/K^+ -ATPase, α_1 and β_1 subunit, mRNA, kidney

Human and experimental diabetes is accompanied by an early, prominent and persistent renal hypertrophy [6, 22]. However, there are some evidence in experimental animal models of diabetes that early insulin treatment can prevent or reverse renal hypertrophy [15]. Nephromegaly has been associated with a renal hyperfunction (increased glomerular filtration rate of Na^+ and water reabsorption,

¹This paper is dedicated to the memory of Professor Tibor Kovács (1929-1994)

Correspondence should be addressed to

Ágota VÉR

Institute of Biochemistry I, Semmelweis University of Medicine

H-1088 Budapest, Puskin u. 9, Hungary