

Available online at www.sciencedirect.com

Biochimica et Biophysica Acta xx (2008) xxx–xxx

www.elsevier.com/locate/bbamcr

Nuclear translocation of the phosphoprotein Hop (Hsp70/Hsp90 organizing protein) occurs under heat shock, and its proposed nuclear localization signal is involved in Hsp90 binding

Sheril Daniel^a, Graeme Bradley^{a,1}, Victoria M. Longshaw^a, Csaba Söti^b, Peter Csermely^b, Gregory L. Blatch^{a,*}

^a Chaperone Research Group, Department of Biochemistry, Microbiology and Biotechnology, Rhodes University, Grahamstown, South Africa

^b Department of Medical Chemistry, Semmelweis University, Puskin street 9, H-1088, Budapest, Hungary

Received 14 September 2007; received in revised form 11 January 2008; accepted 14 January 2008

Abstract

The Hsp70–Hsp90 complex is implicated in the folding and regulation of numerous signaling proteins, and Hop, the Hsp70–Hsp90 Organizing Protein, facilitates the association of this multichaperone machinery. Phosphatase treatment of mouse cell extracts reduced the number of Hop isoforms compared to untreated extracts, providing the first direct evidence that Hop was phosphorylated *in vivo*. Furthermore, surface plasmon resonance spectroscopy (SPR) showed that a cdc2 kinase phosphorylation mimic of Hop had reduced affinity for Hsp90 binding. Hop was predominantly cytoplasmic, but translocated to the nucleus in response to heat shock. A putative bipartite nuclear localization signal (NLS) has been identified within the Hsp90-binding domain of Hop. Although substitution of residues within the major arm of this proposed NLS abolished Hop–Hsp90 interaction as determined by SPR, this was not sufficient to prevent the nuclear accumulation of Hop under leptomycin-B treatment and heat shock conditions. These results showed for the first time that the subcellular localization of Hop was stress regulated and that the major arm of the putative NLS was not directly important for nuclear translocation but was critical for Hop–Hsp90 association *in vitro*. We propose a model in which the association of Hop with Hsp90 and the phosphorylated status of Hop both play a role in the mechanism of nucleocytoplasmic shuttling of Hop.

© 2008 Elsevier B.V. All rights reserved.

Keywords: Hop; Heat shock protein; Phosphorylation; Hsp90; Co-chaperone; Nuclear localization signal

1. Introduction

The Hsp70–Hsp90 Organizing Protein, Hop, was first identified in yeast during a genetic screen for proteins involved in the heat shock response [1]. Hop has insignificant chaperoning abilities [2] and upon heat shock (HS) and viral transformation, no corresponding change in the steady state levels of this protein was detected, although an increase in mRNA levels of Hop in mammalian cells was observed [3,4]. Hop acts as a scaffolding

protein, mediating the interaction of the molecular chaperones Hsp70 and Hsp90 through specific tetratricopeptide-repeat (TPR)-rich binding domains [5,6] to form the Hsp70–Hop–Hsp90 chaperone heterocomplex. These TPR domains characterize the various homologues of Hop, found in humans [3], mice [7], rats [8], insects [9], plants [10], parasites [11] and viruses [12]. Hsp70 binding is mediated primarily through the N-terminal TPR domain of Hop (TPR1) [13] while a central TPR motif-containing region (TPR2A) is known to be essential for Hsp90 binding [14,15]. The TPR-acceptor site on both Hsp70 and Hsp90 comprises an EEVD motif at the C-terminus [15,16].

In the yeast system, it was recently shown that the third TPR domain, TPR2B is also directly involved in Hop's interactions with Hsp70 and Hsp90, and that an isolated TPR2A domain is

* Corresponding author. Tel.: +27 46 603 8262; fax: +27 46 622 3984.

E-mail address: g.blatch@ru.ac.za (G.L. Blatch).

¹ Current Address: Department of Biochemistry and Microbiology, University of Fort Hare, Alice, South Africa.

unable to interact with Hsp90 unless combined with TPR2B [17]. Deletion of a specific amino acid in the TPR2B domain of yeast Hop (A438) showed the disruption of *in vivo* interaction with Hsp90 [18]. The possibility of a network of interactions between Hop and the chaperones Hsp70 and Hsp90, involving domains other than the TPRs, have been suggested, which support its functionality as a scaffolding protein [15,19].

Early studies reported the presence of human Hop in the Golgi apparatus and small vesicles of normal cells, and in the nucleolus of SV40-transformed cells [3]. Immunoprecipitation studies using membranous fractions from mouse brain showed that a population of Hop is localized at the cell surface, where it interacts with Prp^c (a protein whose expression is crucial to the propagation of neurological disease, in particular, Prion disease) both *in vitro* and *in vivo*. It has been proposed that Hop is taken into the plasma membrane as part of a Prp^c–protein complex [20]. Alternatively, Hop may be secreted by a pathway that is distinct from the classical route of membrane translocation through the ER and Golgi apparatus [21] since a transmembrane domain or signal peptide for membrane transport is not found within the amino acid sequence of Hop [4,20].

An investigation into the subcellular localization of Hop in mouse cells showed a predominantly cytoplasmic localization under normal growth conditions [4]. Leptomycin-B treatment (nuclear export inhibition) of mouse fibroblast cells, resulted in a predominantly nuclear accumulation of Hop, suggesting that the protein shuttled between the nucleus and the cytoplasm, with the export of Hop from the nucleus occurring at a faster rate than its import [22]. Moreover, Longshaw et al. identified a potential bipartite nuclear localization signal (NLS), which when fused to enhanced green fluorescent protein (EGFP), resulted in the localization of EGFP to the nucleus [22], providing evidence that this NLS was indeed functional on its own. This putative NLS comprised of a short lysine arm (K222–K223), a 13-amino acid spacer region, followed by a major lysine-rich arm (K237–K239).

Early studies done on yeast Hop [1] identified two major and two minor isoforms of the protein with pIs ranging from 5.75 to 6.05, which implied that Hop was differentially phosphorylated. Other studies showed shifts in the isoform composition of Hop to the more acidic range after viral transformation and heat shock [3,4]. Immediately upstream of the putative bipartite NLS, two potential phosphorylation sites for casein kinase II (CKII; S189) and cell division cycle 2 (*cdc2*; T198) kinase were identified using *in vitro* assays [23]. Despite speculations that Hop is a phosphoprotein, based on indirect evidence, thus far there has been no direct evidence to show that Hop exists as a phosphoprotein *in vivo*.

There is evidence to suggest that the localization of Hop is regulated by phosphorylation. Hydroxyurea (G1/S arrest) and olomoucine treatment (*cdc2* kinase inhibition) of mouse fibroblast cells resulted in cytoplasmic and nuclear localization of Hop [22]. Experiments with phosphorylation mimics of Hop (Hop-S189E and Hop-T198E) suggested that phosphorylation of Hop by *cdc2* kinase promoted the cytoplasmic retention of Hop, whereas phosphorylation of Hop by CKII promoted nuclear localization [22]. Both these putative phosphorylation

sites were found proximal to the Hsp90-binding TPR2A domain of Hop, suggesting that Hop–Hsp90 interactions may also be modulated by phosphorylation. We also noted that the proposed NLS of Hop overlapped with its TPR2A domain. Moreover residues within the spacer region of the NLS (K229, N233 and K301) were shown to be crucial for Hop–Hsp90 interactions [6,15]. Therefore, we hypothesize that the putative bipartite NLS of Hop, and its phosphorylation status, play a significant role in its subcellular localization and association with Hsp90. In this report we have further characterized the putative bipartite NLS of Hop, provided the first direct evidence that Hop was phosphorylated *in vivo*, showed that heat shock affected the subcellular localization and phosphorylation of Hop, and provided evidence that Hop–Hsp90 interactions involved residues in the putative NLS and may be influenced by phosphorylation. 120

2. Experimental procedures 121

2.1. General 122

Common molecular biology procedures such as agarose gel electrophoresis, ligation reactions, competent bacterial cell preparation, plasmid DNA isolation and restriction enzyme digestion were performed according to standard protocols [24]. Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) was according to Laemmli [25], and Western blotting according to Towbin [26]. Luminol-based chemiluminescence detection reagents (ECL Advance Western Blotting Detection Kit; Amersham Biosciences) were used for Western analysis. Protein estimation was done using the 2-D Quant Kit (Amersham Biosciences). 131

2.2. Site-directed mutagenesis 132

All mutations were generated from the plasmids pGEX3X2000 and pSK-mSTII-EGFP, which contain the cDNAs encoding the full-length proteins mSTII (mouse homolog of Hop) as a C-terminal fusion with glutathione-S-transferase (GST) and mSTII as an N-terminal fusion with enhanced green fluorescent protein (EGFP), respectively. All mutations were carried out by site-directed mutagenesis using a double-stranded whole plasmid linear amplification procedure (QuikChange mutagenesis kit; Stratagene). For screening purposes, silent mutations were engineered to create restriction sites except where the desired codon change(s) automatically generated restriction sites. The following mutant proteins were generated: GST-Hop-K239A, GST-Hop/K237/238/239A, GST-Hop-T198E, GST-Hop-S189E, Hop-K239A-EGFP, and Hop-K237/K238/K239A-EGFP. Mutations were confirmed both by restriction enzyme analysis and by DNA sequencing. For ease of manipulation, the mSTII-EGFP encoding fragment in pSK-mSTII-EGFP was used as a template for mutagenesis before subcloning into pBCMGSNeo [27] XhoI/NotI sites to produce pB-mSTII-EGFP and its mutant derivatives, as described previously [22]. Note: All plasmid constructs listed retain their original cloning nomenclature which refers to mSTII (mouse homolog of Hop); however the proteins they encode have been labeled Hop for this study so as to limit the number of acronyms used for the same protein. 151

2.3. Production and purification of GST-Hop fusion proteins 152

Exponentially growing *Escherichia coli* XLI Blue cells carrying pGEX3X153 derived plasmid constructs were induced for 5–6 h at 37 °C with 1 mM isopropyl-1-thio- β -D-galactopyranoside (IPTG). The cells were harvested and lysed by mild sonication in 0.01 culture volume of ice-cold phosphate-buffered saline (PBS; 137 mM NaCl, 2.7 mM KCl, 4.3 mM Na₂HPO₄, 1.4 mM KH₂PO₄, pH 7.3) containing 1 mM final concentration of phenylmethylsulfonyl fluoride (PMSF). The sonicate was incubated, with gentle agitation, for 30 min at room temperature after addition of Triton X-100 to 1% final concentration. The extracts were clarified by centrifugation at 12,000 g for 20 min at 4 °C. Aliquots of clarified extracts were added to 50% (w/v) slurry of glutathione-162

163 agarose beads (2 mL bed volume per 100 mL of extract) previously equi-
 164 librated with PBS. Binding was allowed to occur for 1 h at 4 °C with gentle
 165 rocking. The beads were washed extensively with ice-cold PBS before the
 166 bound GST fusion proteins were eluted by adding appropriate volume of the
 167 elution buffer (10 mM reduced glutathione in 50 mM Tris–HCl, pH 8.0).
 168 Eluted GST fusion proteins were analyzed on SDS-PAGE (12% acrylamide/
 169 bis-acrylamide w/v). GST-Hop and its mutant derivatives were all successfully
 170 expressed and found to be soluble proteins, which were purified to at least 90%
 171 purity.

172 2.4. Surface plasmon resonance spectroscopy

173 Surface plasmon resonance (SPR) spectroscopy was performed using
 174 a Biacore X apparatus (BIAcore, Sweden). A mixture of NHS/EDC
 175 (*N*-hydroxysuccinimide and 1-ethyl-3-[3-dimethylaminopropyl] carbodi-
 176 imide) was used to activate the surface of a CM5 sensor chip by modifica-
 177 tion of the carboxymethyl groups to *N*-Hydroxysuccinimide esters. Purified
 178 rat liver Hsp90 (untagged) was injected into the first flow cell, such that the
 179 *N*-Hydroxysuccinimide esters reacted spontaneously with the amines on the
 180 Hsp90 to form covalent links.

181 The second flow cell was used as a control with no protein immobilized on
 182 it. Approximately 9165 response units (RU) of Hsp90 were immobilized onto
 183 the chip. After equilibration with running buffer (10 mM Hepes buffer con-
 184 taining 140 mM KCl and 0.001% Tween), increasing concentrations (0.2 μ M,
 185 1 μ M and 5 μ M) of GST-Hop and its derivatives were passed over the
 186 immobilized Hsp90 at a flow rate of 10 μ L min⁻¹. GST was passed through both
 187 cells and any non-specific binding to Hsp90 and/or the chip was monitored. All
 188 experiments were performed at room temperature. The CM5 Sensor Chip
 189 (Research Grade) was purchased from BIAcore (Sweden). The data was
 190 analyzed using the BIAevaluation software version 3.0.

191 2.5. Hop isoform analysis by two-dimensional gel electrophoresis

192 To satisfactorily determine the in vivo phosphorylated isoform status of Hop,
 193 NIH 3T3 fibroblast cells were treated in various ways prior to preparation of a
 194 lysate for analysis by two-dimensional (2D) gel electrophoresis. The cells were
 195 either treated with phosphatase inhibitors (PI) okadaic acid (30 nM; specific for
 196 serine/threonine phosphatases) and sodium vanadate (2 mM; specific for tyrosine
 197 phosphatases), to limit or eliminate any dephosphorylation of Hop, or treated

with shrimp alkaline phosphatase (SAP; 10 U) to dephosphorylate any 198
 phosphorylated isoforms of Hop. The NIH 3T3 fibroblast cells treated for 199
 isoform analysis were harvested by scraping into ice-cold PBS. The cell pellets 200
 obtained were resuspended in two-dimensional (2D) lysis buffer (40 mM Tris– 201
 HCl, 8 M Urea, 4% Triton X-100, 1 μ g/mL aprotinin and 100 μ g/mL PMSF) and 202
 lysed on ice for 1 h. The lysate was centrifuged at 13,000 rpm for 30 min on a 203
 benchtop microfuge to remove insoluble cell debris. Protein concentration was 204
 determined using the 2D-Quant kit (Amersham Biosciences, USA), before 205
 the protein samples were precipitated using the 2D clean-up kit (Amersham 206
 Biosciences, USA). 207

For the first dimension separation by isoelectric focusing (IEF), the pre- 208
 cipitated total protein samples were diluted in rehydration buffer (9 M Urea, 2% 209
 Triton X-100, 2% Ampholytes, 0.002% bromophenol blue, 100 mM DTT) to a 210
 final concentration of 1 μ g/ μ L. A total protein sample (120 μ g) was then applied 211
 onto an immobilized linear pH gradient (IPG; pH 4–7) strip within the chambers 212
 of a Zoom® IPGRRunner™ Cassette (Invitrogen) and rehydrated overnight as per 213
 the manufacturer's instructions. Rehydrated IPG strips were focused by per- 214
 forming IEF in distilled water. The applied electric potential (V) was increased in 215
 a step-wise fashion as follows: 200 V (20 min), 450 V (15 min), 750 V (15 min) 216
 and 1000 V (120 min), with a total power of 0.1 W strip⁻¹ and current, 0.05 mA 217
 strip⁻¹. The second dimension was resolved by SDS-PAGE (12% acrylamide/ 218
 bis-acrylamide w/v) at 200 V for 45 min. Western analysis was carried out on the 219
 resolved proteins using antibodies specific to Hop (short chain recombinant 220
 antibodies made to the C-terminal region of mSTI1, C-334; Antibodies By 221
 Design, MorphoSys®; Germany). 222

223 2.6. Cell culture, transfections and preparation of cell lysates

Baby Hamster Kidney (BHK-21) cells and NIH 3T3 mouse fibroblast cells 224
 were maintained in Dulbecco's modified Eagle's medium (Cambrex, USA) 225
 supplemented with 10% calf serum, penicillin (100 U mL⁻¹) and streptomycin 226
 (100 U mL⁻¹) (Highveld Biological PTY. Ltd., South Africa) in a humidified 227
 atmosphere, at 37 °C with 5% CO₂. 228

Transient transfections were conducted on BHK-21 cells using IBAfect (for 229
 magnet assisted transfection) and MA Lipofectin Enhancer (IBA, Germany) on 230
 cells seeded to 60% confluency, according to the manufacturer's instructions. 231
 Leptomycin-B (Synexa Life Sciences, South Africa) was used to arrest the 232
 nuclear export of proteins; cells were treated with 2 ng mL⁻¹ of leptomycin-B 233
 overnight at 37 °C. 234




Fig. 1. Hop was differentially phosphorylated in vivo at 37 °C and 42 °C. (A) The isoform composition of Hop after resolution by 2D gel electrophoresis in the pH range 4–7 of 120 μ g of total protein extract from NIH 3T3 cells (prepared as described in Experimental procedures), followed by Western analysis for the detection of Hop. The isoelectric focusing (IEF) and SDS-PAGE steps are shown. The upper panel shows the Western of lysates prepared from cells that were grown under 37 °C and treated with phosphatase inhibitors (37 °C + PI); the lower panel shows the Western of lysates prepared from cells that were also grown at 37 °C and treated with shrimp alkaline phosphatase (SAP) but not treated with PI (37 °C — PI+SAP). Approximate pIs for the prominent isoforms were estimated and are indicated at the top of each panel. The position of the isoforms with assigned pIs did not always coincide due to technical discrepancies such as slight changes in gel size. The major isoform under normal conditions, with an approximate pI of 6.1 was arbitrarily given the label '0'. Any isoform that was found to be more acidic than '0' was labeled with increasing positive numbers. Additional isoforms which were not assigned a pI value were labeled with an asterisk. This figure was representative of the results obtained from at least three independently conducted experiments. (B). The two panels show cell extracts prepared from cells that were incubated for 30 min at 42 °C. The upper panel shows the isoform composition of Hop in cells lysed in the absence of SAP but presence of PI and the lower panel, cells lysed in the presence of SAP. This figure was representative of the results obtained from at least three independently conducted experiments.





Fig. 2 (continued).

244 (12% acrylamide/bis-acrylamide w/v), and then subjected to Western analysis
 245 using antibodies specific for β -actin (A-2228, Sigma), Hsp72/73 (SPA-
 246 820, StressGen), Hsp40 (Hdj1 homologue; SPA-400, StressGen), Hsp90 α and β
 247 (SPA-835, StressGen), and Hop (Antibodies by Design, MorphoSys®; Germany).

248 3. Results

249 3.1. Hop was phosphorylated in vivo

250 Various different lysates were prepared from NIH 3T3 cells
 251 treated with phosphatase inhibitors (PI) or a phosphatase (SAP).

This allowed the potential phosphorylated state of Hop in vivo to 252
 be analyzed after 2D gel electrophoresis and Western analysis 253
 for the detection of Hop isoforms (Fig. 1A). Previously pub- 254
 lished phosphorylation studies of mouse Hop were performed 255
 using cell lysates that were treated with a tyrosine phosphatase 256
 inhibitor (sodium vanadate) but no serine/threonine phosphatase 257
 inhibitors [4]. As preliminary work to the study presented here, 258
 the 2D-electrophoretic experiments described by Lässle et al. [4] 259
 were repeated but without using any phosphatase inhibitors. The 260
 results were identical to those previously obtained [4] suggesting 261
 that Hop was phosphorylated at sites other than tyrosine. Thus 262

the present study made use of serine/threonine phosphatase inhibitors, in addition to the tyrosine phosphatase inhibitor.

For lysates from cells treated with PI and no SAP added, Hop was found to exist not only as a major and a minor isoform, but also as three more relatively acidic isoforms (Fig. 1A, upper panel). The major isoform with a calculated pI of approximately 6.1 was labeled '0' and the minor isoform found at a pI of approximately 6.0 was labeled '+1'. A third isoform, approaching a pI of 6.0 and merging with isoform +1 was labeled '+2'. The two other relatively more acidic isoforms at pIs less than 5.9 were designated by asterisks and were not assigned approximate pIs. For lysates from cells treated only with SAP there was a considerable reduction in isoforms +1 and +2 (Fig. 1A, lower panel), suggesting that they were dephosphorylated by the SAP treatment. These results provided evidence of *in vivo* phosphorylation of at least two subpopulations of Hop under normal conditions. The most acidic isoform obtained under normal conditions (designated by an asterisk) disappeared after SAP treatment, indicating that this isoform may have corresponded to a minor phosphorylated isoform of Hop. A prominent isoform, designated '+3' was detected after SAP treatment at an approximate pI of 5.9, and this isoform was not prominent in lysate obtained from normally grown cells that were treated with PI (Fig. 1A, upper panel). These results directly related the occurrence of certain acidic isoforms of Hop to post-translational phosphorylation of the protein.

Mouse NIH 3T3 cells were subjected to heat shock, and shown to undergo induction of the stress response as evidenced by induction of Hsp72 protein (data not shown). The isoform composition of Hop after 30 min of heat shock (Fig. 1B, upper panel) differed from that of normal conditions in that there was a decrease in the amount of the major isoform 0. There was also an increase of a number of minor acidic isoforms, the most prominent of which had an approximate pI of 5.9 and was designated '+3'. This isoform was also seen in the SAP treated normal lysate (Fig. 1A, lower panel). The additional minor isoforms that appeared after heat shock were not assigned pIs, and were rather indicated by asterisks. However, it was noted that these isoforms were found at pIs of 5.9 or lower. Lysates prepared after treatment of heat-shocked cells with SAP yielded an isoform profile for Hop (Fig. 1B, lower panel) which included a slight increase of the previously major isoform '0', and the complete absence of isoform +2. Isoform +3 showed a considerable reduction after treatment with SAP. These results were consistent with previous studies [4] that showed that heat shock resulted in modifications of Hop. This data also provided evidence for the first time that Hop was phosphorylated *in vivo* after heat shock.

3.2. The major arm of the proposed bipartite NLS did not contribute to the nuclear localization of Hop under leptomycin-B treatment

We examined the subcellular distribution of Hop in baby hamster kidney cells (BHK-21), using a Hop-EGFP fusion protein, which could also be modified to determine the possible effects of mutations in the predicted NLS. The protein was

transiently expressed in BHK-21 cells, fixed, and then analyzed by confocal laser fluorescence microscopy.

Under normal conditions Hop-EGFP was found to localize predominantly to the cytoplasm of most cells (Fig. 2A, I; B, I; C), whereas the localization pattern of EGFP was found to be distributed throughout the cell and therefore identical to previously published findings [data not shown; [4,22]]. Treatment of transfected cells with leptomycin-B, a specific inhibitor of the CRM-1 nuclear protein export pathway, showed the accumulation of Hop-EGFP in the nucleus (Fig. 2A II; C), a finding that correlates with previous literature [22].

We mutated the residues of the major arm of the proposed NLS, a region in most bipartite NLSs that is critical for functionality *in vivo* [28–30], and examined the subcellular localization patterns of the derivative proteins. Both NLS mutants of Hop-EGFP also showed a predominantly cytoplasmic profile of subcellular localization under normal growth conditions (Fig. 2A, I; B, I). However, closer inspection of the various localization patterns by quantitative analysis revealed that both of the mutant Hop-EGFP derivatives showed a greater incidence of cytoplasmic and nuclear localization compared to unmutated Hop-EGFP (Fig. 2C). Leptomycin-B treated cells showed that both mutant Hop-EGFP derivatives were 'arrested' within the nucleus, as was the case for unmutated Hop-EGFP (Fig. 2A, II; C).

3.3. Hop was localized to the nucleus under heat shock and the nuclear localization of Hop was not dependent on the major arm of the proposed NLS

BHK-21 cells were treated with pre-warmed DMEM and heat shocked at 42 °C for 30 min after transient transfection. In response to heat shock, Hop and its putative NLS mutants showed increased nuclear localization compared to its subcellular localization under normal conditions (Fig. 2B I and II; C).

Total cell extracts obtained from cells grown at 37 °C and cells heat shocked at 42 °C for 30 min, were biochemically fractionated into cytoplasmic, membrane and nuclear fractions. Volume equivalence of each subcellular fraction was loaded proportional to the identical number of cells they were extracted from. The proteins in each fraction were resolved by SDS-PAGE and then subjected to Western blot analysis using antibodies against β -actin, Hsp72/73, Hsp40, Hsp90 and Hop (Fig. 2D). β -actin was found, as expected, in high amounts in the cytoplasmic and membranous extracts of both normal and heat-shocked extracts (Fig. 2D, Panel 1: lanes 1, 2, 3 and 4). No β -actin was detected in the nuclear fraction from cells grown under normal conditions (Fig. 2D, Panel 1: lane 5), suggesting that this fraction was relatively free of contamination from the cytoplasmic fraction. Although generally not nuclear, β -actin is known to reversibly localize to the nucleus under heat shock [31,32] and a faint band was thus visible in the nuclear extract of heat-shocked extracts (Fig. 2D, Panel 1: lane 6). Hsp72/73 was found to be predominantly cytoplasmic and membranous under normal and heat shock conditions (Fig. 2D, Panel 2: lanes 1–4). There was a slight presence of Hsp72/73 in the nuclear fractions before heat shock and a definite nuclear fractionation of Hsp72/73 under heat shock (Fig. 2D, Panel 2: lanes 5–6).




Fig. 3. Proposed NLS residues were critical for Hop–Hsp90 interaction and the phosphorylation of Hop by cdc2 kinase may perturb Hop–Hsp90 interactions. (A) Binding response curves of GST-Hop and its NLS mutants (0.2–5 μ M used, only 5 μ M trace shown for each; GST-Hop-K239A and GST-Hop-K237/238/239A) to Hsp90 (immobilized) were generated using surface plasmon resonance spectroscopy (SPR). (B) Ribbon representation of the structure of TPR2A of Hop (light grey) with the C-terminal MEEVD motif of Hsp90 shown in dark grey (1ELR; [6]). The space filled residues, K237, K238 and K239 form the major arm of the predicted NLS, and these are shown relative to the space filled residue, methionine, which forms part of the pentapeptide MEEVD of Hsp90. This figure was generated using Pymol Molecular Graphics Software [[52]; <http://pymol.sourceforge.net>]. (C) Binding response curves of GST-Hop and its phosphorylation mimics (GST-Hop-S189E, mimic of phosphorylation by CKII kinase; and GST-Hop-T198, mimic of phosphorylation by cdc2 kinase) to Hsp90 using SPR. GST was used as a control for non-specific binding in A and C.

373 The detection of Hsp72/73 in the nuclear fraction after heat
374 shock was consistent with what has been shown previously
375 [33], and these data together with the actin data, suggested that
376 the nuclear fractions had been correctly isolated.

377 The localization of Hsp40 (Hdj1 homologue) was also ana-
378 lyzed, since it is a co-chaperone of Hsp70, and like Hsp70
379 translocates from the cytoplasm to the nucleus under heat shock
380 conditions [34]. Hsp40 was detected in the cytoplasmic frac-
381 tions under normal and heat-shocked conditions, as was ex-
382 pected, with an increased detection in the membranous fraction
383 after heat shock (Fig. 2D, Panel 3: lanes 1–4). Although Hsp40
384 could not be detected in the nuclear fractions obtained from
385 normally grown cells, it was visibly co-fractionated with Hsp70
386 within the nuclear proteins from heat-shocked cells (Fig. 2D,
387 Panel 3: lanes 5–6). Hsp90 was detected predominantly in the
388 cytoplasmic fraction, with an increased detection in the mem-
389 branous fraction after heat shock (Fig. 2D, Panel 4: lanes 1–4).
390 Although previous reports have shown the nuclear translocation
391 of Hsp90 during stress conditions [5,35,36], Hsp90 was not
392 detected in the heat-shocked nuclear fractions obtained in this
393 study (Fig. 2D, Panel 4: lanes 5–6). Hop also showed pre-
394 dominantly cytoplasmic fractionation under normal and heat
395 shock conditions with some detection in the membranous
396 fraction. This increased after heat shock (Fig. 2D, Panel 5: lanes
397 1–4). The detection of Hop was evident in the nuclear fraction
398 of heat-shocked lysate (Fig. 2D, Panel 5: lane 6). Taken together
399 these results demonstrated that a population of Hop localized to
400 the nucleus when the cells were subjected to heat shock, and
401 that mutations of the major arm of the proposed NLS of Hop did
402 not disrupt this nuclear localization.

403 3.4. Proposed NLS residues were critical for Hop–Hsp90 404 interactions and a cdc2 kinase phosphorylation mimic of Hop 405 reduced its affinity for Hsp90

406 SPR was used to investigate Hop–Hsp90 interactions when
407 one or more residues of the major arm of the proposed NLS of
408 Hop were mutated. Hsp90 was immobilized on the sensor chip
409 and GST-Hop constructs were passed over the chip. GST was
410 used as a negative control to monitor any non-specific binding
411 of GST-Hop and its derivatives to Hsp90. Affinity constants in
412 the form of K_D values were determined for the binding of GST-
413 Hop to Hsp90 and found to be 1.4 μ M, consistent with pre-
414 viously published data [6,15]. Both GST-Hop-K239A and GST-
415 Hop-K237/238/239A showed abrogated binding to Hsp90 and
416 were comparable to the negative control GST (Fig. 3A). These
417 results suggested that residues K237, K238 and K239 were
418 critical for the interaction of Hop with Hsp90. A three dimen-
419 sional (3D) representation of the structure of the TPR2A domain
420 of Hop interacting with the C-terminal MEEVD pentapeptide of
421 Hsp90 (Fig. 3B) showed that K237, K238 and K239 potentially
422 made contact with the methionine of the Hsp90 MEEVD motif.

423 We also investigated the effect of phosphorylation at sites
424 S189 and T198 of Hop on its interactions with Hsp90. Phos-
425 phorylation was mimicked at both these sites by substituting the
426 respective serine and threonine residues with an acidic residue,
427 glutamic acid. SPR analysis was conducted to determine the

428 extent of interaction of both these Hop phosphorylation mimics
429 with Hsp90. The binding curves obtained by plotting response
430 units versus time (seconds) showed that the CKII kinase phos-
431 phosphorylation mimic of Hop (GST-Hop-S189E) did not show
432 much deviation from the unmutated GST-Hop (Fig. 3C) where-
433 as the cdc2 kinase phosphorylation mimic of Hop (GST-Hop-
434 T198E) exhibit diminished affinity for Hsp90. K_D values of
435 2.5 μ M and 1 μ M were obtained for GST-Hop-T198E–Hsp90
436 and GST-Hop-S189E–Hsp90 interactions respectively. These
437 values may or may not be of biological significance in vivo.

438 All SPR analyses were repeated independently by immobi-
439 lizing the GST-Hop constructs on a sensor chip and passing
440 over purified Hsp90. Similar results were obtained (data not
441 shown). Taken together, the results from these experiments
442 showed that the major arm of the proposed NLS was crucial for
443 Hop–Hsp90 interactions, and that mimicking phosphorylation
444 by cdc2 kinase reduced the affinity of Hop for Hsp90.

445 4. Discussion

446 This is the first report to have provided direct evidence of Hop
447 being a phosphoprotein under normal and stress conditions, and
448 that the subcellular localization of Hop was primarily stress
449 regulated. We have established that unlike previously reported
450 bipartite NLSs, the major arm of the proposed NLS of Hop was
451 not critical for its nuclear translocation. However these residues
452 were crucial for Hop–Hsp90 interaction, and through this in-
453 teraction, we hypothesize that this region could possibly serve to
454 retain Hop within the cytoplasm. Furthermore, phosphorylation-
455 mimic studies suggested that phosphorylation of Hop at a cdc2
456 kinase site may diminish the affinity of Hop for Hsp90, whilst
457 phosphorylation of Hop at a CKII site may not interfere with
458 Hop–Hsp90 interactions [22].

459 The presence of okadaic acid (serine/threonine phosphatase
460 inhibitor) in the phosphorylation study conducted, suggested
461 that phosphorylation of Hop occurred predominantly at serine or
462 threonine residues. This is further substantiated by evidence of in
463 vitro phosphorylation of Hop at S189 and T198 [22,23]. Heat
464 stress may exert inhibitory and activating effects on various
465 kinases, including Mitogen-activated protein (MAP) kinases,
466 and phosphatases [37,38,39,40]. Hop was found to have a po-
467 tential phosphorylation site for MAPKAP kinase 2 and two
468 potential sites for S6 kinases within its amino acid sequence [4].
469 Hop may thus be phosphorylated by a stress kinase pathway in
470 vivo [4] in response to heat shock. The differentially phos-
471 phosphorylated isoforms of Hop may correspond to subpopulations
472 of Hop, which are localized to varying subcellular compartments
473 during normal and/or stress conditions. Since Hop is known to
474 be predominantly cytoplasmic under normal conditions in this
475 study and in others [4,22], the major ‘0’ isoform of Hop resolved
476 by 2D analysis, may correspond to the cytoplasmic population of
477 Hop under normal conditions.

478 Acute heat shock resulted in increased incidence of cells
479 showing greater translocation of Hop-EGFP to the nucleus (but
480 not nucleolus). This was supported by the detection of endo-
481 genous Hop in the nuclear protein fractions after heat shock.
482 Short-term heat shock in mammalian cells is known to cause the

483 transient arrest of the cell cycle at the G1/S stage [41]. The arrest
 484 of mouse fibroblast cells in the G1/S phase showed increased
 485 nuclear localization of Hop [22]. Since the heat shock
 486 conditions used here also increased the localization of Hop to
 487 the nucleus, these conditions probably promoted G1/S arrest of
 488 the cells. The nuclear localization of Hop may thus be primarily
 489 stress regulated and secondarily regulated by cell cycle events.

490 Although mutations within the major arm of the putative
 491 NLS of Hop did not disrupt the nuclear translocation of the
 492 Hop-EGFP construct, SPR analysis showed that these muta-
 493 tions resulted in minimal interaction of GST-Hop with Hsp90. A
 494 three-dimensional (3D) representation of the Hop TPR2A do-
 495 main showing the NLS residues that were mutated, together
 496 with the C-terminal pentapeptide of Hsp90 (MEEVD motif),
 497 illustrated the proximity of these NLS residues to the meth-
 498 ionine of the Hsp90 peptide (Fig. 3B). One or more of these
 499 residues could be involved in direct interactions with Hsp90.

500 We propose that the minor arm of this putative NLS could be
 501 the functional NLS, acting together with a second putative
 502 monopartite NLS found within amino acids 337–351 (Fig. 4).
 503 Mutation of the major arm of the putative bipartite NLS may
 504 have resulted in a modification of the structure of Hop such that
 505 the putative monopartite NLS was better positioned and ex-
 506 posed to importin binding, thereby facilitating more efficient

nuclear localization than that obtained for unmutated Hop. A
 507 second possibility is that the major arm of the putative bipartite
 508 NLS of Hop might promote the cytoplasmic retention of Hop by
 509 binding to Hsp90, thus explaining why mutation of residues
 510 within Hop that rendered it incapable of interacting with Hsp90
 511 in vitro, allowed for greater nuclear translocation of Hop in
 512 vivo. Hsp90 has previously been reported to sequester the glu-
 513 cocorticoid receptor to the cytoplasm through their interaction
 514 within a complex [42].

515
 516 A subpopulation of endogenous Hop was detected in the
 517 membranous fractions under normal and heat shock conditions.
 518 This finding was consistent with reports of the interaction of
 519 Hop with Prp^c at the plasma membrane [20]. Interestingly, it
 520 was also found that Prp^c interacts with Hop at residues 230–245
 521 [43], which overlaps with the Hsp90 binding site [[15]; and this
 522 study]. Prp^c binding to Hop at this site may act as a mechanism
 523 to improve the efficiency of retaining a subpopulation of Hop at
 524 the plasma membrane. This further supports the proposed role
 525 of Hop–Hsp90 interactions in modulating the subcellular lo-
 526 calization of Hop. Our data also showed the co-fractionation of
 527 Hsp70, Hsp90 and Hsp40, along with Hop, in the membranes
 528 under normal and heat shock conditions. This also suggests that
 529 Hop may associate with one or more of its chaperone partners
 530 within various membranous structures.




Fig. 4. Positioning of the proposed bipartite NLS with respect to TPR2A domain and a second potential NLS. Ribbon representation of the structure of TPR2A of Hop and a flanking C-terminal helix interacting with the C-terminal MEEVD motif of Hsp90 (1ELR; [6]). The residues, K229, N233 and K301 are those that are known to be critical for interaction with Hsp90. Residues denoted with a single asterisk correspond to those that form the major arm of the bipartite NLS (K237, K238 and K239). The residues denoted with a double asterisk, K222 and K223 (dark grey), form part of the minor arm of this NLS, which may be responsible for functionality of the NLS within the protein. A second NLS has been identified within the flanking C-terminal portion of Hop (337–351) shown in dark grey, which may be working in concert with the first predicted NLS. This figure was generated using Pymol Molecular Graphics Software [49]; <http://pymol.sourceforge.net>.

531 Based on the results obtained from this study, together with
 532 what is currently known about Hop particularly regarding its
 533 nucleo-cytoplasmic shuttling properties, we propose a model
 534 detailed in Fig. 5. Within the normal cytoplasmic environment,
 535 Hop equilibrates between an Hsp90-complexed state and an
 536 uncomplexed free state [44,45], and the equilibrium between
 537 the two states may be regulated through phosphorylation of Hop
 538 (Fig. 5, arrow A). Phosphorylation of Hop by cdc2 kinase may
 539 disrupt Hop–Hsp90 interactions as indicated by the SPR analysis.
 540 This frees Hop, but retains it within the cytoplasm [[22];
 541 Fig. 5, arrow C/cdc2?] so that it can interact with other proteins
 542 in complexes that are independent of Hsp90 [46,47].

543 Under normal conditions, Hop–Hsp90 interactions within the
 544 cytoplasm may serve to anchor Hop to this subcellular compart-
 545 ment. The major arm of the proposed NLS of Hop may

546 contribute to the overall localization of Hop by its interaction
 547 with Hsp90. This facilitates the assembly of the Hsp70–Hsp90
 548 chaperone complex, which has been shown to be cytoplasmic
 549 (Fig. 5, arrow B). It also implies that Hop translocates into the
 550 nucleus when it is not complexed with Hsp90, possibly pro-
 551 moted by CKII phosphorylation of Hop (Fig. 5, arrow C/CKII?).
 552 Previous reports have shown the nuclear translocation of Hsp90
 553 during stress conditions including heat shock [5,35,36]. In the
 554 present study Hsp90 was not detected in the nuclear fractions
 555 under normal or heat shock conditions. Biochemical fractiona-
 556 tion techniques are known to be disruptive and may have altered
 557 the subcellular localization pattern of a protein such as Hsp90,
 558 whose translocation into the nucleus may be transient or dyn-
 559 amic in nature. This may also explain why relatively low levels
 560 of Hop were detected in the nuclear fractions, when compared to




Fig. 5. A model reflecting the possible links between the subcellular localization of the phosphoprotein Hop and its functionality as a Hsp70/Hsp90 scaffolding protein. Hop exists on its own or in complex with Hsp90, in the cytoplasm under normal conditions. This may be regulated by phosphorylation, with cdc2 kinase phosphorylation of Hop disrupting its interaction with Hsp90 (cdc2? arrow; A). Interaction of Hop with Hsp90 is known to facilitate a number of other interactions, of which the most well established one is the interaction of the Hop–Hsp90 complex with Hsp70 (associated with its co-chaperone Hsp40 and substrate) in order to facilitate substrate transfer from Hsp70 to Hsp90 (B). This multichaperone complex then dissociates, freeing its various components. Hop is known to translocate to the nucleus (C) under stressful conditions, and its localization may be regulated by phosphorylation, with CKII phosphorylation possibly promoting nuclear localization (CKII? arrow) and cdc2 kinase phosphorylation possibly promoting cytoplasmic retention (cdc2? arrow). It is speculated that Hop may also be capable of moving into the nucleus in concert with Hsp90 (D) as a complex (arrows shown in dotted lines) by either the putative NLS (222–239), or through the functioning of multiple NLSs, and possibly also promoted by CKII phosphorylation (CKII? dotted arrow). It is already known that both Hsp70 (together with Hsp40) and Hsp90 translocate into the nucleus under heat shock (E and F respectively). Within the nucleus Hop may have a number of functions, including its basic function of interacting with Hsp70 and/or Hsp90 to form nuclear complexes (G). The 1, 2A, 2B, C and NLS annotations on Hop refer to its TPR1, TPR2A and TPR2B domains, C-terminal domain, and nuclear localization signal sequence, respectively. Hsp40, Hsp70 and Hsp90 are labeled as 40, 70 and 90, respectively.

the confocal microscopic data. Thus, if Hsp90 does indeed translocate into the nucleus during heat shock, the absence of functional NLSs within Hsp90 [36,48] suggests that Hsp90 may have to “piggyback” on an NLS-bearing protein. CKII phosphorylation was shown to promote nuclear localization of Hop [22] and now, evidence has been presented to suggest that CKII phosphorylation does not disrupt Hop–Hsp90 binding. Thus we cannot exclude the possibility of a subpopulation of Hop–Hsp90 translocating as a complex into the nucleus under specific regulatory conditions, such as phosphorylation by CKII (Fig. 5, dotted arrow D/CKII?). As other studies [34] and ours have shown the presence of Hsp70 and Hsp40 in heat-shocked nuclear fractions. This is suggestive of the nuclear assembly of the Hsp70/Hsp90 chaperone heterocomplex during heat stress (Fig. 5 arrows E, F and G).

The details of the co-localization of Hop and Hsp90, as well as the scaffolding properties of Hop within the nucleus, under conditions of heat shock remains to be investigated. Changes in the ionic state and conformation of Hop due to phosphorylation at S189 and T198, may have an effect on its interactions with other members of the chaperone heterocomplex, such as Hsp70, and this also needs to be investigated. The mechanism of nuclear localization of the Hsp70–Hop–Hsp90 chaperone heterocomplex under both normal and stress conditions is complex. However, the putative phosphorylation-regulated NLS of Hop appears to play an important role in the functionality of the chaperone heterocomplex, not only on the level of its subcellular localization but also with regards to the assembly of the chaperone heterocomplex.

Acknowledgements

This work was funded in part by a National Research Foundation (NRF) Unlocking the Future Focus Area grant (South Africa; GUN No: 2053542) and a Wellcome Trust grant (UK; Grant No: 066705) awarded to GLB, and a NRF–Hungary Collaborative grant (South Africa; GUN No: 2067467) awarded to GLB and PC. SD was awarded an Andrew Mellon Scholarship (Rhodes University, South Africa) during her doctoral studies, and a JRC Adhoc Postdoctoral Bursary (Rhodes University, South Africa) to facilitate the writing of this paper. CS is a Bolyai Research Scholar of the Hungarian Academy of Sciences. The authors thank Bronwyn McLean (Graphics Services Unit, Rhodes University, South Africa) for her expert assistance with the figures presented. We also thank the staff of the Electron Microscope Unit at University of Kwazulu-Natal, Pietermaritzburg (South Africa) for their technical assistance.

References

- [1] C.M. Nicolet, E.A. Craig, Isolation and characterization of STI1, a stress-inducible gene from *Saccharomyces cerevisiae*, *Mol. Cell. Biol.* 9 (1989) 3638–3646.
- [2] R.J. Schumacher, R. Hurstll, W.P. Sullivan, N.J. McMahons, D.O. Tofts, R.L. Matts, ATP-dependent chaperoning activity of reticulocyte lysate, *J. Biol. Chem.* 269 (1994) 9493–9499.
- [3] B. Honoré, H. Leffers, P. Madsen, H.H. Rasmussen, J. Vandekerckhove, J.E. Celis, Molecular cloning and expression of a transformation-sensitive human

- protein containing the TRP motif and sharing identity to the stress-inducible yeast protein STI1, *J. Biol. Chem.* 267 (1992) 8485–8491.
- [4] M. Lässle, G.L. Blatch, V. Kundra, T. Takatori, B.R. Zetter, Stress-inducible murine protein mSTI1. Characterization of binding domains for heat shock proteins and in vitro phosphorylation by different kinases, *J. Biol. Chem.* 272 (1997) 1876–1884.
- [5] W.B. Pratt, D.O. Toft, Steroid receptor interactions with heat shock protein and immunophilin chaperones, *Endocr. Rev.* 18 (1997) 306–360.
- [6] C. Scheufler, A. Brinker, G. Bourenkov, S. Pegoraro, L. Moroder, H. Bartunik, F.U. Hartl, I. Moarefi, Structure of TPR domain–peptide complexes: critical elements in the assembly of the Hsp70–Hsp90 multi-chaperone machine, *Cell* 101 (2000) 199–210.
- [7] G.L. Blatch, M. Lässle, B.R. Zetter, V. Kundra, Isolation of a mouse cDNA encoding mSTI1, a stress-inducible protein containing the TPR motif, *Gene* 194 (1997) 277–282.
- [8] J. Demand, J. Luders, J. Hohfeld, The carboxy-terminal domain of Hsc70 provides binding sites for a distinct set of chaperone cofactors, *Mol. Cell. Biol.* 18 (1998) 2023–2028.
- [9] M.D. Adams, S.E. Celniker, R.A. Holt, C.A. Evans, J.D. Gocayne, P.G. Amanatides, S.E. Scherer, P.W. Li, R.A. Hoskins, R.F. Galle, et al., The genome sequence of *Drosophila melanogaster*, *Science* 287 (2000) 2185–2195.
- [10] Z. Zhang, M.K. Quick, K.C. Kanelakis, M. Gijzen, P. Krishna, Characterization of a plant homolog of Hop, a cochaperone of Hsp90, *Plant Physiol.* 131 (2003) 525–535.
- [11] J.R. Webb, A. Campos-Neto, Y.A.W. Skeiky, S.G. Reed, Molecular characterization of the heat-inducible LmSTI1 protein of *Leishmania major*, *Mol. Biochem. Parasitol.* 89 (1997) 179–193.
- [12] C.H. Cheng, S.M. Liu, T.Y. Chow, Y.Y. Hsiao, D.P. Wang, J.J. Huang, H. H. Chen, Analysis of the complete genome sequence of the Hz-1 virus suggests that it is related to members of the Baculoviridae, *J. Virol.* 76 (2002) 9024–9034.
- [13] J. van der Spuy, B.D. Kana, H.W. Dirr, G.L. Blatch, Heat shock cognate protein 70 chaperone-binding site in the co-chaperone murine stress-inducible protein 1 maps to within three consecutive tetratricopeptide repeat motifs, *Biochem. J.* 345 (2000) 645–651.
- [14] S. Chen, V. Prapapanich, R.A. Rimerman, B. Honore, D.F. Smith, Interactions of p60, a mediator of progesterone receptor assembly, with heat shock proteins Hsp90 and Hsp70, *Mol. Endocrinol.* 10 (1996) 682–693.
- [15] O.O. Odunuga, J.A. Hornby, C. Bies, R. Zimmermann, D.J. Pugh, G.L. Blatch, Tetratricopeptide repeat motif-mediated Hsc70–mSTI1 interaction. Molecular characterization of the critical contacts for successful binding and specificity, *J. Biol. Chem.* 278 (2003) 6896–6904.
- [16] A. Brinker, C. Scheufler, F. von der Mulbe, B. Fleckenstein, C. Herrmann, G. Jung, I. Moarefi, F.U. Hartl, Ligand discrimination by TPR domains. Relevance and selectivity of EEVD-recognition in hsp70.hsp90 complexes, *J. Biol. Chem.* 277 (2002) 19265–19275.
- [17] G. Flom, R.H. Behal, L. Rosen, D.G. Cole, J.L. Johnson, Definition of the minimal fragments of Sti1 required for dimerization, interaction with Hsp70 and Hsp90 and in vivo functions, *Biochem. J.* 404 (2007) 158–167.
- [18] G. Flom, J. Weekes, J.J. Williams, J.L. Johnson, Effect of mutation of the tetratricopeptide repeat and aspartate-proline 2 domains of Sti1 on Hsp90 signaling and interaction in *Saccharomyces cerevisiae*, *Genetics* 172 (2006) 41–51.
- [19] P.E. Carrigan, D.L. Riggs, M. Chinkers, D.F. Smith, Functional comparison of human and drosophila Hop reveals novel role in steroid receptor maturation, *J. Biol. Chem.* 280 (2005) 8906–8911.
- [20] S.M. Zanata, M.H. Lopes, A.F. Mercadante, G.N.M. Hajj, L.B. Chiarini, R. Nomizo, A.R.O. Freitas, A.L.B. Cabral, K.S. Lee, M.A. Juliano, E. de Oliveira, S.G. Jachieri, R. Linden, R.R. Brentani, V.R. Martins, Stress-inducible protein 1 is a cell surface ligand for cellular prion that triggers neuroprotection, *EMBO J.* 21 (2002) 3307–3316.
- [21] A. Muesch, E. Hartmann, K. Rohde, R. Rubartelli, R. Sitia, T.A. Rapoport, A novel pathway for secretory proteins? *Trends. Biochem. Sci.* 15 (1990) 86–88.
- [22] V.M. Longshaw, J.P. Chapple, M.S. Balda, M.E. Cheetham, G.L. Blatch, Nuclear translocation of the Hsp70/Hsp90 organizing protein mSTI1 is regulated by cell cycle kinases, *J. Cell. Sci.* 117 (2004) 701–710.

- 683 [23] V.M. Longshaw, H.W. Dirr, G.L. Blatch, M. Lässle, The in vitro phospho-
684 phosphorylation of the co-chaperone mSTII by cell cycle kinases substantiates
685 a predicted casein kinase II-p34cdc2-NLS (CeN) motif, *Biol. Chem.* 381
686 (2000) 1133–1138.
- 687 [24] J. Sambrook, D.W. Russell, *Molecular Cloning: a Laboratory Manual*, 3
688 ed., Cold Spring Harbour Laboratory Press, New York, 2001.
- 689 [25] U.K. Laemmli, Cleavage of structural proteins during the assembly of the
690 head of bacteriophage T4, *Nature* 227 (1970) 680–685.
- 691 [26] H. Towbin, T. Staehelin, J. Gordon, Electrophoretic transfer of proteins
692 from polyacrylamide gels to nitrocellulose sheets: procedure and some
693 applications, *Proc. Natl. Acad. Sci. U. S. A.* 76 (1979) 4350–4354.
- 694 [27] H. Karasuyama, F. Melchers, Establishment of mouse cells which constitu-
695 tively secrete large quantities of interleukin 2, 3, 4, or 5, using modified
696 cDNA expression vectors, *Eur. J. Immunol.* 18 (1988) 97–104.
- 697 [28] E. Conti, J. Kuriyan, Crystallographic analysis of the specific yet versatile
698 recognition of distinct nuclear localization signals by karyopherin alpha,
699 *Structure* 8 (2000) 329–338.
- 700 [29] M.R. Fontes, T. The, B. Kobe, Structural basis of recognition of monopartite
701 and bipartite nuclear localization sequences by mammalian importin-alpha,
702 *J. Mol. Biol.* 297 (2000) 1183–1194.
- 703 [30] B. Friedrich, C. Quensel, T. Sommer, E. Hartmann, M. Köhler, Nuclear
704 localization signal and protein context mediate importin a specificity of
705 nuclear import substrates, *Mol. Cell. Biol.* 26 (2006) 8697–8709.
- 706 [31] K. Iida, S. Matsumoto, I. Yahara, The KKRKK Sequence is involved in
707 heat shock-induced nuclear translocation of the 18-kDa actin-binding
708 protein, Cofilin, *Cell Struct. Funct.* 17 (1992) 39–46.
- 709 [32] A. Wada, M. Fukuda, M. Mishima, E. Nishida, Nuclear export of actin: a
710 novel mechanism regulating the subcellular localization of a major cyto-
711 skeletal protein, *EMBO J.* 17 (1998) 1635–1641.
- 712 [33] J.M. Velazquez, S. Lindquist, Hsp70: nuclear concentration during envi-
713 ronmental stress and cytoplasmic storage during recovery, *Cell* 36 (1984)
714 655–662.
- 715 [34] H. Hattori, Y-C. Liu, I. Tohna, M. Ueda, T. Kaneda, T. Kobayashi, K.
716 Tanabe, K. Ohtsuka, Intracellular localization and partial amino acid
717 sequence of a stress-inducible 40-kDa protein in HeLa cells, *Cell Struct.*
718 *Funct.* 17 (1992) 77–86.
- 719 [35] M. Biggiogera, R.M. Tanguay, R. Marin, Y. Wu, T.E. Martin, S. Fakan,
720 Localization of heat shock proteins in mouse male germ cells: an immu-
721 noelectron microscopical study, *Exp. Cell Res.* 229 (1996) 77–85.
- 722 [36] T. Langer, S. Rosmus, H. Fasold, Intracellular localization of the 90 kDa
723 heat shock protein (HSP90alpha) determined by expression of a EGFP-
724 HSP90alpha-fusion protein in unstressed and heat stressed 3T3 cells, *Cell*
725 *Biol. Int.* 27 (2003) 47–52.
- [37] G. Bagi, E.J. Hidvegi, Protein phosphorylation and kinase activities in 726
tumour cells after hyperthermia, *Int. J. Radiat. Biol.* 58 (1990) 633–650. 727
- [38] G.N. Somero, Proteins and temperature, *Annu. Rev. Physiol.* 57 (1995) 728
43–68. 729
- [39] J. Rouse, P. Cohen, S. Trigon, M. Morange, A. Alonso-Llamazares, A 730
novel kinase cascade triggered by chemical stress and heat shock which 731
stimulates MAP kinase-activated protein kinase-2 and phosphorylation of 732
the small heat shock proteins, *Cell* 78 (1994) 1027–1037. 733
- [40] F. Chen, M. Torres, R.F. Duncan, Activation of mitogen-activated protein 734
kinase by heat shock treatment in *Drosophila*, *Biochem. J.* 312 (1995) 735
341–349. 736
- [41] E.W. Khandijan, Heat treatment induces dephosphorylation of pRb and 737
dissociation of T-antigen pRb complex during transforming infection with 738
SV40, *Oncogene* 10 (1995) 359–367. 739
- [42] A. Guiochon-Mantel, K. Delabre, P. Lescop, E. Milgrom, Intracellular 740
traffic of steroid hormone receptors, *J. Steroid Biochem. Mol. Biol.* 56 741
(1996) 3–9. 742
- [43] A.S. Coitinho, M.H. Lopes, G.N.M. Hajj, J.I. Rossato, A.R. Freitas, C.C. 743
Castro, M. Cammarota, R.R. Brentani, I. Izquierdo, V.R. Martins, Short- 744
term memory formation and long-term memory consolidation are enhanced 745
by cellular prion association to a stress-inducible protein 1, *Neurobiol. Dis.* 746
26 (2007) 282–290. 747
- [44] H.C.J. Chang, D.F. Nathan, S. Lindquist, In vivo analysis of the hsp90 748
cochaperone sti1 (p60), *Mol. Cell. Biol.* 17 (1997) 318–325. 749
- [45] M.P. Hernandez, W.P. Sullivan, D.O. Toft, The assembly and intermo- 750
lecular properties of the hsp70–Hop–hsp90 molecular chaperone complex, 751
J. Biol. Chem. 277 (2002) 38294–38304. 752
- [46] T. Abbas-Terki, O. Donze, P.A. Briand, D. Picard, Hsp104 interacts with Hsp90 753
cochaperones in respiring yeast, *Mol. Cell. Biol.* 21 (2001) 7569–7575. 754
- [47] M. Gebauer, R. Melki, U. Gehring, The chaperone cofactor hop/p60 in- 755
teracts with the cytosolic chaperonin-containing TCP-1 and affects its nu- 756
cleotide exchange and protein folding activities, *J. Biol. Chem.* 273 (1998) 757
29475–29480. 758
- [48] K.I.L. Kang, J. Devin, F. Cadepond, N. Jibard, A. Guiochon-Mantel, E.-M. 759
Baulieu, M.G. Catelli, In vivo functional protein–protein interaction: 760
nuclear targeted Hsp90 shifts cytoplasmic steroid receptor mutants into the 761
nucleus, *Proc. Natl. Acad. Sci. U. S. A.* 91 (1994) 340–344. 762
- [49] W.L. DeLano, S. Bromberg, *Pymol Users Guide*, DeLano Scientific LLC, 763
California, 2004. 764